

Letter to the Editor

Regarding Trisomy 2p Syndrome

To the Editor:

We read with interest the publication by Hahm et al. [1999], "Trisomy 2p syndrome: A fetus with anencephaly and postaxial polydactyly." We would like to report on another case of anencephaly in a fetus with triplication of 2p23.1→2pter resulting from a paternally derived translocation. The karyotype in this case was 46,XX,der(3)t(2;3)(p23.1;q29)pat.

The fetus was diagnosed at 15 weeks with anencephaly when the mother presented for amniocentesis on account of the known paternal translocation. Autopsy confirmed anencephaly and demonstrated, in addition, a sacral meningomyelocoele; no other abnormalities were detected.

A paternal sister with the same unbalanced translocation is now 25 years old. She was not available for examination but is said to be severely retarded with absent tear ducts and a scoliosis. She is able to walk.

Neural tube defects have now been reported in several cases with trisomy 2p [Fineman et al., 1983; Sarda et al., 1992; Singer and Gerson, 1987; Thirkelsen et al., 1973; Winsor et al., 1997]. The triplicated segments consisted of 2p23→2pter, 2p13→2pter, 2p24→2pter, 2p13→2pter, and 2p13→2p24, respectively. As in case and that of Hahm et al., all included 2p24.

The case presented here adds further support to the suggestion by Lurie et al. [1995] and Winsor et al.

[1997] that 2p24 is a candidate region for genes responsible for neural tube development.

REFERENCES

- Fineman RM, Buyse M, Morgan M. 1983. Variable phenotype associated with duplication of different regions of 2p. *Am J Med Genet* 15:451–456.
- Hahm GK, Barth RF, Schauer GM, Reiss R, Opitz JM. 1999. Trisomy 2p syndrome: a fetus with anencephaly and postaxial polydactyly. *Am J Med Genet* 87:45–48.
- Lurie IW, Ilyina HG, Gurevich DB, Rumyantseva NV, Naumchik IV, Castellani C, Hoeller A, Schinzel A. 1995. Trisomy 2p: analysis of unusual phenotypic findings. *J Med Genet* 55:229–236.
- Sarda P, Lefort G, Devaux P, Humeau C, Rieu D. 1992. Multiple congenital anomalies due to partial 2p13→2pter duplication resulting from an unbalanced X;2 translocation. *Ann Génét* 35:117–120.
- Singer N, Gersen S. 1987. The value of chromosome analysis in cases of neural tube defects: a case of anencephaly associated with fetal dup(2)(p24→pter). *Prenat Diagn* 7:567–571.
- Thirkelsen AJ, Hultén M, Jonasson J, Lindsten J, Christensen NC, Iversen T. 1973. Presumptive direct insertion within chromosome 2 in man. *Ann Hum Genet* 36:367–373.
- Winsor S, McGrath M, Khalifa M, Duncan A. 1997. A report of recurrent anencephaly with trisomy 2p23-2pter: additional evidence for the involvement of 2p24 in neural tube development and evaluation of the role for cytogenetic analysis. *Pren Diagn* 17:665–669.

Diana Wellesley*

Wessex Clinical Genetics Service
Princess Anne Hospital
Southampton, United Kingdom

Tracy Boyle

Wessex Regional Genetics Laboratory
Salisbury District Hospital
Salisbury, United Kingdom

*Correspondence to: Dr. Diana Wellesley, Wessex Clinical Genetics Service, Princess Anne Hospital, Coxford Road, Southampton SO16 5YA, U.K.

Received 3 January 2000; Accepted 14 February 2000